


PERCUTANEOUS VERTEBROPLASTY AND BALLOON KYPHOPLASTY

SUPPLEMENT



PERCUTANEOUS VERTEBROPLASTY AND BALLOON KYPHOPLASTY **SUPPLEMENT**

CHRIS DE LAET, NANCY THIRY, KIRSTEN HOLDT HENNINGSSEN, SABINE STORDEUR, CÉCILE CAMBERLIN



COLOPHON

Title :	Percutaneous vertebroplasty and balloon kyphoplasty – Supplement
Authors :	Chris De Laet (KCE), Nancy Thiry (KCE), Kirsten Holdt Henningsen (KCE), Sabine Stordeur (KCE), Cécile Camberlin (KCE)
Project coordinator and Senior supervisor:	Sabine Stordeur (KCE)
Reviewers :	Frank Hulstaert (KCE), Raf Mertens (KCE), Lorena San Miguel (KCE)
External Experts :	Michael Bruneau (Belgian Society of Neurosurgery (BSN) – Hôpital Erasme, Bruxelles), Philippe Claesen (Jessa Ziekenhuis), Geert Crombez (UGent), Bart Depreitere (UZ Leuven), Hendrik Fransen (AZ St-Lucas Gent), Patrick Galloo (Socialistische Mutualiteiten), Alphonse Lubansu (Hôpital Erasme Bruxelles), Germain Milbouw (CHR Namur), Henri Nielens (Cliniques universitaires Saint-Luc, Bruxelles), Valérie Noblesse (INAMI – RIZIV), Bart Poffyn (UZ Gent), Stéphane Sobczak (AXXON), Johan Van Lerbeirghe (SSBE Spine Society of Belgium), Jan Van Meirhaeghe (AZ St-Jan Brugge), Patrick Van Schaeybroeck (Imelda Ziekenhuis, Bonheiden), Peter Van Wambeke (UZ Leuven), Dominique Verhulst (ZNA Stuivenberg, Antwerpen), René Westhovens (UZ Leuven)
External Validators :	Wilco Jacobs (The Health Scientist, The Netherlands), Christian Raftopoulos (Cliniques universitaires St-Luc, Belgium), Matt Stevenson (University of Sheffield, The United Kingdom)
Acknowledgements:	UNAMEC (Fédération belge de l'industrie des technologies médicales – Belgische federatie van de industrie van de medische technologiën), Nicolas Fairon (KCE): information specialist
Other reported interests:	<p>Membership of a stakeholder group on which the results of this report could have an impact: Wilco Jacobs (member of various focused spinal surgery associations), Johan Van Lerbeirghe (SSBE), Michael Bruneau (Hôpital Erasme – Université Libre de Bruxelles)</p> <p>Owner of subscribed capital, options, shares or other financial instruments: Wilco Jacobs (Clinical Research consultancy The Health Scientist)</p> <p>Fees or other compensation for writing a publication or participating in its development: Wilco Jacobs (Scientific collaborator for systematic literature research about lumbar disc prostheses)</p> <p>Participation in scientific or experimental research as an initiator, principal investigator or researcher: Wilco Jacobs (PI ZonMW funds with cofinancing from Medtronic Inc for minimal invasive lumbar fusion, Principal Investigator for systematic literature research about spine); Bart Poffyn (Head Researcher 'MISS Anterior Approach Tumors fractures in spine')</p> <p>Grants, fees or funds for a member of staff or another form of compensation for the execution of research: Wilco Jacobs (Fees from LUMC for collaboration to the guideline development about robot spinal surgery)</p> <p>Consultancy or employment for a company, an association or an organisation that may gain or lose financially</p>



due to the results of this report: Dominique Verhulst (DePuy Spine (Johnson & Johnson)
Payments to speak, training remuneration, subsidised travel or payment for participation at a conference: Christian Raftopoulos (Johnson & Johnson conferences); Jan Van Meirhaeghe
Presidency or accountable function within an institution, association, department or other entity on which the results of this report could have an impact: Johan Van Lerbeirghe (president SSBC); Patrick Galloo (President Implants and Invasive Medical Devices Reimbursement Commission); Patrick Van Schaeybroeck (Vice-President Spine Society Belgium; BNSS, Board Member Belgian Neurosurgical Spine Society)

Layout : Ine Verhulst, Joyce Grijseels

Disclaimer :

- The external experts were consulted about a (preliminary) version of the scientific report. Their comments were discussed during meetings. They did not co-author the scientific report and did not necessarily agree with its content.
- Subsequently, a (final) version was submitted to the validators. The validation of the report results from a consensus or a voting process between the validators. The validators did not co-author the scientific report and did not necessarily all three agree with its content.
- Finally, this report has been approved by common assent by the Executive Board.
- Only the KCE is responsible for errors or omissions that could persist. The policy recommendations are also under the full responsibility of the KCE

Publication date

29 October 2015

Domain:

Health Technology Assessment (HTA)

MeSH :

Vertebroplasty; Kyphoplasty; Fractures, Compression

NLM Classification :

WE 725

Language :

English

Format :

Adobe® PDF™ (A4)

Legal depot :

D/2015/10.273/101

ISSN:

2466-6459

Copyright :

KCE reports are published under a "by/nc/nd" Creative Commons Licence
<http://kce.fgov.be/content/about-copyrights-for-kce-reports>.



How to refer to this document ?

De Laet C, Thiry N, Holdt Henningsen K, Stordeur S, Camberlin C. Percutaneous vertebroplasty and balloon kyphoplasty – Supplement. Health Technology Assessment (HTA) Brussels: Belgian Health Care Knowledge Centre (KCE). 2015. KCE Reports 255. D/2015/10.273/101.

This document is available on the website of the Belgian Health Care Knowledge Centre



■ APPENDIX REPORT

TABLE OF CONTENTS

■	APPENDIX REPORT	1
1.	BELGIAN SITUATION.....	2
1.1.	REIMBURSEMENT SCHEDULE	2
1.1.1.	INAMI – RIZIV billing codes for vertebroplasty and balloon kyphoplasty	2
1.1.2.	Reimbursement conditions for balloon kyphoplasty:	3
1.2.	DIAGNOSES FOUND IN THE RHM-MZG DATA	4
1.2.1.	Most frequent 3-digit ICD-9-CM codes of principal diagnosis in the case of vertebroplasty ...	4
1.2.2.	Five-digit ICD-9-CM codes of principal diagnosis in the case of Other disorders of bone and cartilage (733.xx) in the case of vertebroplasty	4
1.2.3.	Most frequent 3-digit ICD-9-CM codes of principal diagnosis in the case of kyphoplasty.....	5
1.2.4.	Five-digit ICD-9-CM codes of principal diagnosis in the case of Other disorders of bone and cartilage (733.xx) in the case of kyphoplasty	5
2.	CLINICAL EFFECTIVENESS AND SAFETY.....	6
2.1.	PICO	6
2.1.1.	PICO for Vertebroplasty and selection criteria.....	6
2.1.2.	PICO for Kyphoplasty and selection criteria	7
2.2.	SEARCH STRATEGY	8
2.2.1.	General search strategy.....	8
2.2.2.	Detailed search strategy	8
2.3.	SEARCH RESULTS.....	12
2.3.1.	Flow chart of selection for systematic search	13
2.4.	METHODOLOGICAL QUALITY.....	14
2.4.1.	Systematic review	14
2.4.3.	RCTs	15
3.	LITERATURE SEARCH FOR ECONOMIC EVALUATIONS.....	16
3.1.	SEARCH STRATEGY	16
3.2.	SELECTION PROCEDURE	18
■	REFERENCES	20



1. BELGIAN SITUATION

1.1. Reimbursement schedule

1.1.1. INAMI – RIZIV billing codes for vertebroplasty and balloon kyphoplasty

Code	Label (Fr / NL)	Key letter	Tariff (*)
Procedure for balloon kyphoplasty			
(589676) - 589680	Cyphoplastie percutanée par ballonnet pour le traitement de fractures par enfoncement d'un corps vertébral (par corps vertébral) / Percutane ballonkyphoplastie voor de behandeling van indeukingsfracturen van een wervellichaam (per wervellichaam)	I 400	€ 305.35
Balloon kyphoplasty material before July the 1st, 2014			
(683012) - 683023	Matériel de consommation pour cyphoplastie avec ballon, utilisé à l'occasion de la prestation 589676-589680/ Gebruiksmateriaal voor kyphoplastie met ballon gebruikt naar aanleiding van de verstrekking 589676-589680	U 4121	
(683034) - 683045	Ciment pour cyphoplastie avec ballon utilisé conjointement à la prestation 683012-683023, par niveau, maximum 2 niveaux / Cement voor kyphoplastie met ballon gebruikt samen met de verstrekking 683012-683023, per niveau, maximum 2 niveaus	U 82	
Balloon kyphoplasty material between July the 1st, 2014 and April the 1st, 2015.			
(162971) - 162982	Matériel de consommation pour cyphoplastie avec ballon, utilisé à l'occasion de la prestation 589676-589680 de la nomenclature / Gebruiksmateriaal voor kyphoplastie met ballon gebruikt naar aanleiding van de verstrekking 589676-589680 van de nomenclatuur		€ 4257.64
(162993) - 163004	Ciment pour cyphoplastie avec ballon utilisé conjointement à la prestation 162971-162982, par niveau, maximum deux niveaux / Cement voor kyphoplastie met ballon gebruikt samen met de verstrekking 162971-162982, per niveau, maximum twee niveaus		€ 82.08
Balloon kyphoplasty and vertebroplasty material from April the 1st, 2015.			
(162993) - 163004	Ciment pour cyphoplastie avec ballon utilisé conjointement à la prestation 162971-162982, par niveau, maximum deux niveaux / Cement voor kyphoplastie met ballon gebruikt samen met de verstrekking 162971-162982, per niveau, maximum twee niveaus		€ 82.08
(171975) - 171986	Ciment pour vertébroplastie, par niveau, maximum 2 niveaux / Cement voor vertebroplastie, per niveau, maximum 2 niveaus		€ 82.08

(Ambulatory) - (*) situation at 01/08/2015

1.1.2. Reimbursement conditions for balloon kyphoplasty:

Reimbursement is limited to

Vertebral compression fractures caused by osteoporosis, single- or multi-level, inferior to T5, without neurological compression and without neurological defect sign, if the following criteria are met:

- Persistent pain of more than 8 weeks linked to the fracture and for which the pain medication therapy administered during at least 8 weeks has failed or caused very troublesome side effects; the situation must have been written in a report by the physician in charge of the pain treatment; the intervention must take place in the 4 months after the first objective findings of the VCF unless a progression of the fracture has been demonstrated.
- Loss of height between 20% and 60%.
- Dorsal wall is maintained.
- Presence of a bony oedema.
- Fractures classified A1.1, A1.2 , A1.3 or A2.3 according to the AO classification (Arbeitsgemeinschaft für Osteosynthesefragen classification).
- In a man with a T-score, calculated in comparison with a male reference population, during a dual energy xray absorptiometry examination:
 - <-2.5 at the lumbar level (L1-L4 or L2-L4) OR
 - <-1 at the hip level (total or femoral neck zone).
- In a woman with a glucocorticoid-induced osteoporosis, treated since more than 3 months with a daily oral dose of 7.5 mg prednisone or equivalent, administered in a recognized scientific indication:
 - The patient must be post-menopausal without hormonal substitution OR

- she must present a T-score <-1.5 at the level of the lumbar spine (L1-L4 or L2-L4) or at the hip level (total or femoral neck zone), calculated in comparison with a female reference population, during a dual energy xray absorptiometry examination.

- In a post-menopausal woman presenting a a T-score <-2.5 at the level of the lumbar spine (L1-L4 or L2-L4) or at the hip level (total or femoral neck zone), calculated in comparison with a female reference population, during a dual energy xray absorptiometry examination.

B. Vertebral compression fractures caused by Kahler's myeloma without neurological compression and without neurological defect sign, if the following criteria are met:

- Persistent pain of more than 6 weeks linked to the fracture and for which the pain medication therapy administered during at least 6 weeks has failed or caused very troublesome side effects; the situation must have been written in a report by the physician in charge of the pain treatment.
- Dorsal wall is maintained.
- Presence of a bony oedema.
- Fractures classified A1.1, A1.2., A1.3 or A2.3 according to the AO classification (Arbeitsgemeinschaft für Osteosynthesefragen classification).

In both cases (A and B), this diagnosis is made based on the three following examinations:

- An antero-posterior plain film radiography; and
- a MRI or CT-scan when there is contraindication for MRI; and
- a bone scintigraphy without SPECT; and
- a DEXA-scan in case of osteoporotic VCF.

The documents proving that the conditions are satisfied must be kept in a file and provided to the advisory physician if requested.

A maximum of 2 BKPs can be billed per two calendar years (exceptions can be made requiring the authorization of the College of Medical Directors).



1.2. Diagnoses found in the RHM-MZG data

1.2.1. Most frequent 3-digit ICD-9-CM codes of principal diagnosis in the case of vertebroplasty

Ranking	3 digit ICD-9-CM code	Number of stays	Percentage
1	733 OTHER DISORDERS OF BONE AND CARTILAGE	1107	51.90%
2	805 FRACTURE OF VERTEBRAL COLUMN WITHOUT MENTION OF SPINAL CORD INJURY	566	26.53%
3	722 INTERVERTEBRAL DISC DISORDERS	111	5.20%
4	198 SECONDARY MALIGNANT NEOPLASM - METASTASE - OF OTHER SPECIFIED SITES	72	3.38%
5	724 OTHER AND UNSPECIFIED DISORDERS OF BACK	28	1.31%
6	203 MULTIPLE MYELOMA AND IMMUNOPROLIFERATIVE NEOPLASMS	24	1.13%
7	721 SPONDYLOSIS AND ALLIED DISORDERS	24	1.13%
8	996 COMPLICATIONS PECULIAR TO CERTAIN SPECIFIED PROCEDURES	22	1.03%
9	228 HEMANGIOMA AND LYMPHANGIOMA, ANY SITE	20	0.94%
10	491 CHRONIC BRONCHITIS	14	0.66%
11	Other	145	6.8%
TOTAL		2133	100%

Source: RHM-MZG 2008-2011

1.2.2. Five-digit ICD-9-CM codes of principal diagnosis in the case of Other disorders of bone and cartilage (733.xx) in the case of vertebroplasty

Ranking	5 digit ICD-9-CM code	Number of stays	Percentage
1	733.00 OSTEOPOROSIS, UNSPECIFIED	10	0.47%
2	733.01 SENILE OSTEOPOROSIS	2	0.09%
3	733.13 PATHOLOGIC FRACTURE OF VERTEBRAE	1048	49.13%
4	733.19 PATHOLOGIC FRACTURE OF OTHER SPECIFIED SITE	2	0.09%
5	733.22 ANEURYSMAL BONE CYST	3	0.14%
6	733.49 ASEPTIC NECROSIS OF BONE, OTHER	1	0.05%
7	733.81 MALUNION OF FRACTURE	2	0.09%
8	733.82 NONUNION OF FRACTURE	27	1.27%
TOTAL		1107	51.90%

Source: RHM-MZG data 2008-2011



1.2.3. Most frequent 3-digit ICD-9-CM codes of principal diagnosis in the case of kyphoplasty

Ranking	3 digit ICD-9-CM code	Number of stays	Percentage
1	733 OTHER DISORDERS OF BONE AND CARTILAGE	1507	64.08%
2	805 FRACTURE OF VERTEBRAL COLUMN WITHOUT MENTION OF SPINAL CORD INJURY	610	25.94%
3	203 MULTIPLE MYELOMA AND IMMUNOPROLIFERATIVE NEOPLASMS	24	1.02%
4	724 OTHER AND UNSPECIFIED DISORDERS OF BACK	24	1.02%
5	198 SECONDARY MALIGNANT NEOPLASM - METASTASE - OF OTHER SPECIFIED SITES	21	0.89%
6	722 INTERVERTEBRAL DISC DISORDERS	15	0.64%
7	491 CHRONIC BRONCHITIS	11	0.47%
8	338 PAIN, NOT ELSEWHERE CLASSIFIED	10	0.43%
9	737 CURVATURE OF SPINE	10	0.43%
10	806 FRACTURE OF VERTEBRAL COLUMN WITH SPINAL CORD INJURY	9	0.38%
11	Other	111	4.72%
TOTAL		2352	100%

Source: RHM-MZG 2008-2011

1.2.4. Five-digit ICD-9-CM codes of principal diagnosis in the case of Other disorders of bone and cartilage (733.xx) in the case of kyphoplasty

Ranking	5 digit ICD-9-CM code	Number of stays	Percentage
1	733.00 OSTEOPOROSIS, UNSPECIFIED	21	0.89%
2	733.01 SENILE OSTEOPOROSIS	4	0.17%
3	733.09 OTHER OSTEOPOROSIS	5	0.21%
4	733.10 PATHOLOGIC FRACTURE, UNSPECIFIED SITE	3	0.13%
5	733.13 PATHOLOGIC FRACTURE OF VERTEBRAE	1455	61.86%
6	733.19 PATHOLOGIC FRACTURE OF OTHER SPECIFIED SITE	3	0.13%
7	733.20 CYST OF BONE (LOCALIZED), UNSPECIFIED	1	0.04%
8	733.81 MALUNION OF FRACTURE	2	0.09%
9	733.82 NONUNION OF FRACTURE	12	0.51%
10	733.95 STRESS FRACTURE OF OTHER BONE	1	0.04%
TOTAL		1507	64.01%

Source: RHM-MZG data 2008-2011



2. CLINICAL EFFECTIVENESS AND SAFETY

2.1. PICO

2.1.1. PICO for Vertebroplasty and selection criteria

Selection criteria	Inclusion criteria	Exclusion criteria
Population	Adult patients (18 years of age or older) with vertebral compression fracture due to osteoporosis, malignancy (metastases, myeloma), haemangioma, osteonecrosis or trauma.	
Intervention	Vertebroplasty	
Comparator	<ul style="list-style-type: none">• Conservative (non-surgical) treatment only for osteoporotic VCF or combined with specific radiation therapy, chemotherapy or surgical tumour removal for VCF due to other causes• Kyphoplasty	
Outcomes	<ul style="list-style-type: none">• Mortality• Pain and analgesics use• Mobility• Functional capacity• Health Related Quality of Life• Spine anatomy• Safety	
Study design	HTAs, systematic reviews, randomised controlled trials	Narrative reviews, observational studies, letters, editorials, notes, abstracts



2.1.2. PICO for Kyphoplasty and selection criteria

Selection criteria	Inclusion criteria	Exclusion criteria
Population	Adult patients (18 years of age or older) with vertebral compression fracture due to osteoporosis, malignancy (metastases, myeloma), haemangioma, osteonecrosis or trauma.	
Intervention	Kyphoplasty	
Comparator	<ul style="list-style-type: none">• Conservative (non-surgical) treatment only for osteoporotic VCF or combined with specific radiation therapy, chemotherapy or surgical tumour removal for VCF due to other causes• Vertebroplasty	
Outcomes	<ul style="list-style-type: none">• Mortality• Pain and analgesics use• Mobility• Functional capacity• Health Related Quality of Life• Spine anatomy• Safety	
Study design	HTAs, systematic reviews, randomised controlled trials	Narrative reviews, observational studies, letters, editorials, notes, conference abstracts



2.2. Search strategy

2.2.1. General search strategy

The search is primarily based on the MeSH term Vertebroplasty introduced in 2008. Additionally, a new MeSH term Kyphoplasty was introduced in 2011 but this term is included within the Vertebroplasty term and was therefore not added as a separate MeSH term in the search. However, additional search terms, both as MeSH and free text were added to allow for indexing errors or missing indexing.

Additionally, we consulted the EUneHTA POP^a and INAHTA databases and the ClinicalTrials.gov database (search words ‘vertebroplasty’ OR ‘kyphoplasty’).

2.2.2. Detailed search strategy

Medline RCT (OVID)

ID	Search	Hits
1	vertebra.ab,ti.	9132
2	vertebrae.ab,ti.	14 594
3	cervical vertebrae/	28 150
4	Lumbar vertebrae/	40 134
5	Spine/	23 500
6	Low Back Pain/	14 942
7	1 or 2 or 3 or 4 or 5 or 6	110 983
8	Fractures, Compression/	1274
9	(fracture? adj3 compression).ab,ti.	3697
10	8 or 9	4115
11	(fracture? adj3 (spin* or vertebr*)).ab,ti.	12 787
12	Spinal fracture/	10 640
13	11 or 12	17 931
14	7 and 10	1675

15	13 or 14	18 199
16	cement*.ab,ti.	38 628
17	balloon?.ab,ti.	50 781
18	leakage.ab,ti.	43 648
19	polymethylmethacrylate.ab,ti.	3270
20	pmma.ab,ti.	6547
21	Polymethyl Methacrylate/	5440
22	16 or 17 or 18 or 19 or 20 or 21	139 683
23	15 and 22	1598
24	exp vertebroplasty/	1545
25	vertebroplast*.ab,ti.	2263
26	kyphoplast*.ab,ti.	1135
27	24 or 25 or 26	3073
28	23 or 27	3280
29	limit 28 to yr="2006 -Current"	2544
30	limit 29 to systematic reviews	119
31	randomized controlled trial.pt.	400 323
32	controlled clinical trial.pt.	90 694
33	randomized.ti,ab.	341 710
34	placebo.ti,ab.	168 242
35	clinical trials as topic/	176 382
36	randomly.ti,ab.	229 371
37	trials.ti.	50 693
38	31 or 32 or 33 or 34 or 35 or 36 or 37	949 689
39	exp animal/ not humans/	4 094 649
40	38 not 39	876 116
41	29 and 40	257
42	41 not 30	205

^a POP: Planned and Ongoing Projects Database of projects by EUneHTA members

**Medline SR (OVID)**

ID	Search	Hits
1	vertebra.ab,ti.	9122
2	vertebrae.ab,ti.	14 572
3	cervical vertebrae/	28 135
4	Lumbar vertebrae/	40 105
5	Spine/	23 487
6	Low Back Pain/	14 927
7	1 or 2 or 3 or 4 or 5 or 6	110 899
8	Fractures, Compression/	1272
9	(fracture? adj3 compression).ab,ti.	3693
10	8 or 9	4111
11	(fracture? adj3 (spin* or vertebr*)).ab,ti.	12 782
12	Spinal fracture/	10 632
13	11 or 12	17 922
14	7 and 10	1671
15	13 or 14	18 190
16	cement*.ab,ti.	38 621
17	balloon?.ab,ti.	50 773
18	leakage.ab,ti.	43 611
19	polymethylmethacrylate.ab,ti.	3269
20	pmma.ab,ti.	6543
21	Polymethyl Methacrylate/	5435
22	16 or 17 or 18 or 19 or 20 or 21	139 629
23	15 and 22	1595
24	exp vertebroplasty/	1539
25	vertebroplast*.ab,ti.	2260
26	kyphoplast*.ab,ti.	1134
27	24 or 25 or 26	3070
28	23 or 27	3277
29	(malign* or neoplasm* or tumo?r* or	2 272 687

metasta* or myelom* or haemangiom* or cancer*).ab,ti.

30	exp neoplasms/	2 703 544
31	osteopor*.ab,ti.	54 449
32	exp Osteoporosis/	46 502
33	osteonecro*.ab,ti.	6933
34	exp Osteonecrosis/	12 406
35	trauma*.ab,ti.	258 855
36	exp "Wounds and Injuries"/	737 109
37	29 or 30 or 31 or 32 or 33 or 34 or 35 or 36	4 092 763
38	28 and 37	2878
39	28 not 38	399
40	limit 28 to yr="2006 -Current"	2541
41	limit 40 to systematic reviews	119

EMBASE RCT

ID	Search	Hits
#1	vertebra:ab,ti	11 742
#2	vertebrae:ab,ti	17 843
#3	'spine'/exp	133 859
#4	'low back pain'/exp	37 180
#5	#1 OR #2 OR #3 OR #4	172 653
#6	'compression fracture'/exp	3133
#7	(fracture* NEAR/3 compression):ab,ti	4928
#8	#6 OR #7	6008
#9	(fracture* NEAR/3 (spin* OR vertebr*)):ab,ti	17 402
#10	'spine fracture'/exp	23 447
#11	#9 OR #10	28 831
#12	#5 AND #8	2726
#13	#11 OR #12	29 008
#14	cement*:ab,ti	41 854
#15	balloon*:ab,ti	73 592



#16	leakage*:ab,ti	53 874
#17	polymethylmethacrylate:ab,ti	3529
#18	pmma:ab,ti	7097
#19	'poly(methyl methacrylate)'/exp	11 070
#20	#14 OR #15 OR #16 OR #17 OR #18 OR #19	178 258
#21	#13 AND #20	2359
#22	'percutaneous vertebroplasty'/exp	4413
#23	vertebroplast*:ab,ti	3161
#24	kyphoplast*:ab,ti	1714
#25	#22 OR #23 OR #24	4893
#26	#21 OR #25	5208
#27	malign*:ab,ti OR neoplasm*:ab,ti OR tumor*:ab,ti OR tumour*:ab,ti OR metasta*:ab,ti OR myelom*:ab,ti OR haemangioma*:ab,ti OR cancer*:ab,ti	2 773 585
#28	'neoplasm'/exp	3 461 911
#29	osteoporo*:ab,ti	76 879
#30	'osteoporosis'/exp	94 330
#31	osteonecro*:ab,ti	8467
#32	'bone necrosis'/exp	21 995
#33	trauma*:ab,ti	323 443
#34	'injury'/exp	1 585 751
#35	#27 OR #28 OR #29 OR #30 OR #31 OR #32 OR #33 OR #34	5 503 461
#36	#26 AND #35	4659
#37	#36 AND (2006:py OR 2007:py OR 2008:py OR 2009:py OR 2010:py OR 2011:py OR 2012:py OR 2013:py OR 2014:py)	3774
#38	#26 AND (2006:py OR 2007:py OR 2008:py OR 2009:py OR 2010:py OR 2011:py OR 2012:py OR 2013:py OR 2014:py)	4215
#39	[medline]/lim	21 490 853
#40	#38 NOT #39	1862

#41	[cochrane review]/lim OR 'systematic review' OR 'meta analyse' OR [meta analysis]/lim OR [systematic review]/lim OR 'meta analyses' OR 'meta analysis' OR 'guideline' OR 'guidelines'	600 691
#42	#40 AND #41	106
#43	#37 NOT #39	1696
#44	#43 AND #41	100
#45	#42 NOT #44	6
#46	random*:ab,ti OR placebo*:de,ab,ti OR (double NEXT/1 blind*):ab,ti	1 141 517
#47	#40 AND #46	234

EMBASE SR

ID	Search	Hits
#1	vertebra:ab,ti	11 742
#2	vertebrae:ab,ti	17 843
#3	'spine'/exp	133 859
#4	'low back pain'/exp	37 180
#5	#1 OR #2 OR #3 OR #4	172 653
#6	'compression fracture'/exp	3133
#7	(fracture* NEAR/3 compression):ab,ti	4928
#8	#6 OR #7	6008
#9	(fracture* NEAR/3 (spin* OR vertebr*)):ab,ti	17 402
#10	'spine fracture'/exp	23 447
#11	#9 OR #10	28 831
#12	#5 AND #8	2726
#13	#11 OR #12	29 008
#14	cement*:ab,ti	41 854
#15	balloon*:ab,ti	73 592
#16	leakage*:ab,ti	53 874
#17	polymethylmethacrylate:ab,ti	3529
#18	pmma:ab,ti	7097

#19	'poly(methyl methacrylate)'/exp	11 070
#20	#14 OR #15 OR #16 OR #17 OR #18 OR #19	178 258
#21	#13 AND #20	2359
#22	'percutaneous vertebroplasty'/exp	4413
#23	vertebroplast*:ab,ti	3161
#24	kyphoplast*:ab,ti	1714
#25	#22 OR #23 OR #24	4893
#26	#21 OR #25	5208
#27	malign*:ab,ti OR neoplasm*:ab,ti OR tumor*:ab,ti OR tumour*:ab,ti OR metasta*:ab,ti OR myelom*:ab,ti OR haemangioma*:ab,ti OR cancer*:ab,ti	2 773 585
#28	'neoplasm'/exp	3 461 911
#29	osteoporo*:ab,ti	76 879
#30	'osteoporosis'/exp	94 330
#31	osteonecro*:ab,ti	8467
#32	'bone necrosis'/exp	21 995
#33	trauma*:ab,ti	323 443
#34	'injury'/exp	1 585 751
#35	#27 OR #28 OR #29 OR #30 OR #31 OR #32 OR #33 OR #34	5 503 461
#36	#26 AND #35	4659
#37	#36 AND (2006:py OR 2007:py OR 2008:py OR 2009:py OR 2010:py OR 2011:py OR 2012:py OR 2013:py OR 2014:py)	3774
#38	#26 AND (2006:py OR 2007:py OR 2008:py OR 2009:py OR 2010:py OR 2011:py OR 2012:py OR 2013:py OR 2014:py)	4215
#39	[medline]/lim	21 490 853
#40	#38 NOT #39	1862
#41	[cochrane review]/lim OR 'systematic review' OR 'meta analyse' OR [meta analysis]/lim OR [systematic review]/lim OR 'meta	600 691

	analyses' OR 'meta analysis' OR 'guideline' OR 'guidelines'
#42	#40 AND #41
#43	#37 NOT #39
#44	#43 AND #41
#45	#42 NOT #44

Cochrane

ID	Search	Hits
#1	vertebra:ab,ti	526
#2	vertebrae:ab,ti	526
#3	MeSH descriptor: [Cervical Vertebrae] explode all trees	766
#4	MeSH descriptor: [Lumbar Vertebrae] explode all trees	2130
#5	MeSH descriptor: [Spine] this term only	606
#6	MeSH descriptor: [Low Back Pain] this term only	2100
#7	#1 or #2 or #3 or #4 or #5 or #6	5526
#8	MeSH descriptor: [Fractures, Compression] this term only	95
#9	(fracture* near/3 compression):ab,ti	263
#10	#8 or #9	280
#11	(fracture* near/3 (spin* or vertebr*)):ab,ti	1215
#12	MeSH descriptor: [Spinal Fractures] this term only	630
#13	#11 or #12	1341
#14	#7 and #10	70
#15	#13 or #14	1349
#16	cement*:ab,ti	2246
#17	balloon*:ab,ti	3041
#18	leakage:ab,ti	1899



#19	polymethylmethacrylate:ab,ti	155
#20	pmma:ab,ti	307
#21	MeSH descriptor: [Polymethyl Methacrylate] explode all trees	176
#22	#16 or #17 or #18 or #19 or #20 or #21	7426
#23	#15 and #22	123
#24	MeSH descriptor: [Vertebroplasty] explode all trees	119
#25	vertebroplast*:ab,ti	160
#26	kyphoplast*:ab,ti	126
#27	#24 or #25 or #26	249
#28	#23 or #27	255
#29	#28 Publication Year from 2006 to 2014	219


2.3. Search results

From the systematic search in bibliographical databases we originally retrieved 766 references (including many duplicates). From these we selected and reviewed in detail thirteen recent (since 2010) HTAs and systematic reviews. Thirteen RCTs (seventeen publications) were retrieved in full text. We also selected two meta-analyses. Most reviews included only evidence on osteoporotic VCF but there were two on VCF due to malignancies.

The search through the EUnetHTA POP database and the INAHTA database showed no current new projects, but showed two finished projects with HTA reports published in 2014 (one unpublished at the moment of the search). The additional retrieval of clinical trials through the clinicaltrials.gov database yielded 51 clinical trials, the majority within the scope of our PICO. Those are not listed in the flow chart. While writing this report additional full-text articles were retrieved when needed.



2.3.1. Flow chart of selection for systematic search



SR: systematic review, RCT: randomised controlled trial, MA: meta-analysis.



2.4. Methodological quality

Selected studies were evaluated on their methodological quality using the Cochrane's risk of bias tool for RCTs,¹ and the AMSTAR checklist for systematic reviews.²

2.4.1. Systematic review

AMSTAR checklist of systematic reviews for selected SR

	ScHARR ³
Was an 'a priori' design provided?	Yes
Was there duplicate study selection and data extraction?	Partly*
Was a comprehensive literature search performed?	Yes
Was the status of publication an exclusion criterion?	Yes
Was a list of studies (included and excluded) provided?	Partly
Were the characteristics of the included studies provided?	Yes
Was the scientific quality of the included studies assessed and documented?	Yes
Was the scientific quality of the included studies used appropriately in formulating conclusions?	Yes
Were the methods used to combine the findings of studies appropriate?	Yes
Was the likelihood of publication bias assessed?	No
Was the conflict of interest stated? No Yes	Yes
Total	8/11

*Partly: 10% sample checked by second reviewer, data extraction by two reviewers



2.4.3. RCTs

Cochrane risk of bias table (+= low risk, -= high risk, ?=unclear risk)

RCT	Buchbinder	INVEST	VERTOS	Rousing	VERTOS II	Farrokhi	Blasco	FREE	Liu	KAVIAR	CAFE*
Random sequence generation (selection bias)	+	+	+	+	+	+	?	+	+	+	+
Allocation concealment (selection bias)	+	+	+	+	+	+	?	+	?	?	?
Blinding of participants and personnel (performance bias)	+	+	-	-	-	-	-	-	?	?	-
Blinding of outcome assessment (detection bias)	+	+	-	-	-	-	-	-	+	?*	-
Incomplete outcome data (attrition bias)	+	+	?	?	?	+	?	?	?	?	?
Selective reporting (reporting bias)	?	-	?	?	+	?	+	-	?	?	?
Other bias	+	+	?	?	-	+	?	+	?	?	?

*Blinded central radiologist for radiologic outcomes

The Cochrane risk of bias table (except for the KAVIAR and CAFE trials) is based on the SCHARR report that assesses the internal and external validity of the included trials. The report concludes: “The quality of the included studies is generally not very high. Much of this is owing to the widespread lack of blinding: the studies at least risk of bias are the double-blinded Buchbinder et al.⁴ and INVEST⁵ studies, which compare vertebroplasty with an OPLA. The studies which compare vertebroplasty with non-invasive management (Blasco et al.,⁶ Farrokhi et al.,⁷ Rousing et al.,^{8, 9} VERTOS¹⁰ and VERTOS II¹¹) vary in quality, that by Farrokhi et al. being at least risk of bias.

Only three studies (Blasco et al.,⁶ FREE^{12, 13} and INVEST⁵) appeared to be adequately powered for at least their primary outcomes (pain score in Blasco et al. and INVEST, change in SF-36 PCS score from baseline to 1 month in FREE). Because most studies were underpowered for most outcomes, the absence of a statistically significant treatment effect should not necessarily be taken as evidence that no such difference exists.¹³

The quality of the studies on the effectiveness and safety of vertebroplasty and kyphoplasty for the treatment of non-osteoporotic VCF is very low. Most studies are not-randomised while the only RCT identified (CAFE) is small, allowed for cross-over and is industry sponsored. Obviously, setting up trials for those patients is more time- and money intensive because there are fewer patients than for osteoporotic VCF.



3. LITERATURE SEARCH FOR ECONOMIC EVALUATIONS

3.1. Search strategy

Database			Ovid MEDLINE(R) In-Process & Other Non-Indexed Citations and Ovid MEDLINE(R) 1946 to Present
Date	20/04/2015		
Date covered	1946 to present		
Search strategy	#	Searches	Results
	1	Economics/	26 593
	2	"Costs and Cost Analysis"/	42 278
	3	"Value of Life"/ec [Economics]	227
	4	exp Economics, Pharmaceutical/ or Economics, Medical/ or Economics, Hospital/ or Economics, Dental/ or Economics, Nursing/	26 722
	5	budget\$.ti,ab.	20 160
	6	cost-benefit.mp.	66 258
	7	cost-effectiveness.mp.	37 970
	8	cost-utility.mp.	2753
	9	(cost-minimisation or cost-minimization).mp.	907
	10	or/1-9	187 214
	11	limit 10 to letter	6075
	12	limit 10 to editorial	5092
	13	limit 10 to historical article	2480
	14	or/11-13	13 604

15	10 not 14	173 610
16	Animals/	5 442 465
17	15 not 16	167 274
18	Cervical Vertebrae/	27 933
19	Lumbar Vertebrae/	40 004
20	Spine/	23 300
21	Low Back Pain/	14 831
22	18 or 19 or 20 or 21	98 974
23	Fractures, Compression/	1288
24	(fracture* adj1 compression).ab,ti.	2911
25	23 or 24	3356
26	22 and 25	1158
27	exp Vertebroplasty/	1581
28	vertebroplast*.ab,ti.	2294
29	kyphoplast*.ab,ti.	1159
30	27 or 28 or 29	3129
31	26 or 30	3798
32	limit 31 to (editorial or historical article or letter)	205
33	31 not 32	3593
34	17 and 33	41
35	limit 34 to yr="2006 -Current"	35

Note mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier.

Database	Embase	
Date	20/04/2015	
Date covered	No restriction	
Search strategy	# Searches	Results
#1	vertebra:ab,ti	12 246
#2	vertebrae:ab,ti	18 481
#3	'spine':exp	140 413
#4	'low back pain':exp	38 397
#5	#1 OR #2 OR #3 OR #4	180 382
#6	'compression fracture':exp	3354
#7	(fracture* NEAR/1 compression):ab,ti	4037
#8	#6 OR #7	5222
#9	(fracture* NEAR/1 (spin* OR vertebr*)):ab,ti	12 671
#10	'spine fracture':exp	24 356
#11	#9 OR #10	27 196
#12	#5 AND #8	2727
#13	#11 OR #12	27 519
#14	cement*:ab,ti	42 938
#15	balloon*:ab,ti	76 048
#16	leakage*:ab,ti	55 884
#17	polymethylmethacrylate:ab,ti	3615
#18	pmma:ab,ti	7333
#19	'poly(methyl methacrylate)':exp	11 409
#20	#14 OR #15 OR #16 OR #17 OR #18 OR #19	183 985

#21	#13 AND #20	2345
#22	'percutaneous vertebroplasty':exp	4618
#23	vertebroplast*:ab,ti	3295
#24	kyphoplast*:ab,ti	1806
#25	#22 OR #23 OR #24	5112
#26	#21 OR #25	5402
#27	'cost benefit analysis':exp	66 597
#28	'cost effectiveness analysis':exp	104 949
#29	'cost utility analysis':exp	6001
#30	'cost minimization analysis':exp	2630
#31	'pharmacoeconomics':exp	170 072
#32	#27 OR #28 OR #29 OR #30 OR #31	316 674
#33	#26 AND #32	172
#34	#33 AND ([editorial]/lim OR [letter]/lim OR [note]/lim)	24
#35	#33 NOT #34	148
#36	#35 AND [2006-2015]/py	126



Database CRD HTA and CRD NHS EED		
Date	20/04/2015	
Date covered		No restriction
Search strategy	#	Searches Results
1	(vertebra:ab,ti)	0
2	(vertebrae:ab,ti)	0
3	(MeSH descriptor: [Cervical Vertebrae] explode all trees)	59
4	(MeSH descriptor: [Lumbar Vertebrae] explode all trees)	172
5	(MeSH descriptor: [Spine] this term only)	29
6	(MeSH descriptor: [Low Back Pain] this term only)	208
7	(#1 or #2 or #3 or #4 or #5 or #6)	424
8	(MeSH descriptor: [Fractures, Compression] this term only)	22
9	((fracture* near/3 compression):ab,ti)	0
10	(#8 or #9)	22
11	((fracture* near/3 vertebr*):ab,ti)	0
12	(MeSH descriptor: [Spinal Fractures] this term only)	89
13	#11 OR #12	89
14	#7 AND #10	3
15	#13 OR #14	91
16	(cement*:ab,ti)	17
17	(balloon*:ab,ti)	115
18	(leakage:ab,ti)	0
19	(polymethylmethacrylate:ab,ti)	0

20	(pmma:ab,ti)	1
21	(MeSH descriptor: [Polymethyl Methacrylate] explode all trees)	8
22	(#16 or #17 or #18 or #19 or #20 or #21)	138
23	(#15 and #22)	16
24	(MeSH descriptor: [Vertebroplasty] explode all trees)	34
25	(vertebroplast*:ab,ti)	40
26	(kyphoplast*:ab,ti)	33
27	(#24 or #25 or #26)	60
28	(#23 or #27)	60
29	(#28) FROM 2006 TO 2015	42


Note Searches limited to the HTA and NHS EED databases

3.2. Selection procedure

The electronic searches returned 203 citations in total (35 in Medline(OVID), 126 in Embase and 42 in CRD HTA & CRD NHS EED). No other publication was identified via manual search. After exclusion of 33 duplicates, 170 unique citations were left. The flow chart of the selection process is presented below.



Flowchart of the economic literature review process





■ REFERENCES

1. Higgins JP, Altman DG, Gotzsche PC, Juni P, Moher D, Oxman AD, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ*. 2011;343:d5928.
2. Shea BJ, Hamel C, Wells GA, Bouter LM, Kristjansson E, Grimshaw J, et al. AMSTAR is a reliable and valid measurement tool to assess the methodological quality of systematic reviews. *J Clin Epidemiol*. 2009;62(10):1013-20.
3. Stevenson M, Gomersall T, Jones M.L, Rawdin A, Hernandez M, Dias S, et al. Percutaneous vertebroplasty and percutaneous balloon kyphoplasty for the treatment of osteoporotic vertebral fractures: A systematic review and cost-effectiveness analysis. 2014. *Health Technol. Assess.* 1817 Available from: http://www.journalslibrary.nihr.ac.uk/_data/assets/pdf_file/0017/114317/FullReport-hta18170.pdf
4. Buchbinder R, Osborne RH, Ebeling PR, Wark JD, Mitchell P, Wriedt C, et al. A randomized trial of vertebroplasty for painful osteoporotic vertebral fractures. *The New England journal of medicine*. 2009;361(6):557-68.
5. Kallmes DF, Comstock BA, Heagerty PJ, Turner JA, Wilson DJ, Diamond TH, et al. A randomized trial of vertebroplasty for osteoporotic spinal fractures.[Erratum appears in N Engl J Med. 2012 Mar 8;366(10):970]. *N Engl J Med*. 2009;361(6):569-79.
6. Blasco J, Martinez-Ferrer A, Macho J, San Roman L, Pomes J, Carrasco J, et al. Effect of vertebroplasty on pain relief, quality of life, and the incidence of new vertebral fractures: a 12-month randomized follow-up, controlled trial. *J Bone Miner Res*. 2012;27(5):1159-66.
7. Farrokhi MR, Alibai E, Maghami Z. Randomized controlled trial of percutaneous vertebroplasty versus optimal medical management for the relief of pain and disability in acute osteoporotic vertebral compression fractures. *Journal of neurosurgery. Spine*. 2011;14(5):561-9.



8. Rousing R, Hansen KL, Andersen MO, Jespersen SM, Thomsen K, Lauritsen JM. Twelve-months follow-up in forty-nine patients with acute/semiacute osteoporotic vertebral fractures treated conservatively or with percutaneous vertebroplasty: a clinical randomized study. *Spine*. 2010;35(5):478-82.
9. Rousing R, Andersen MO, Jespersen SM, Thomsen K, Lauritsen J. Percutaneous vertebroplasty compared to conservative treatment in patients with painful acute or subacute osteoporotic vertebral fractures: three-months follow-up in a clinical randomized study. *Spine*. 2009;34(13):1349-54.
10. Voormolen MH, Mali WP, Lohle PN, Fransen H, Lampmann LE, Graaf Y, et al. Percutaneous vertebroplasty compared with optimal pain medication treatment: short-term clinical outcome of patients with subacute or chronic painful osteoporotic vertebral compression fractures. The VERTOS study. *AJNR Am J Neuroradiol*. 2007;28(3):555-60.
11. Klazen CA, Lohle PN, de Vries J, Jansen FH, Tielbeek AV, Blonk MC, et al. Vertebroplasty versus conservative treatment in acute osteoporotic vertebral compression fractures (Vertos II): an open-label randomised trial. *Lancet*. 2010;376(9746):1085-92.
12. Wardlaw D, Cummings SR, Meirhaeghe J, Bastian L, Tillman JB, Ranstam J, et al. Efficacy and safety of balloon kyphoplasty compared with non-surgical care for vertebral compression fracture (FREE): a randomised controlled trial. *Lancet*. 2009;373(9668):1016-24.
13. Boonen S, Meirhaeghe J, Bastian L, Cummings SR, Ranstam J, Tillman JB, et al. Balloon kyphoplasty for the treatment of acute vertebral compression fractures: 2-year results from a randomized trial. *Journal of bone and mineral research*. 2011;26(7):1627-37.